

Sensitive dependence and dense periodic points

S.Kanmani

Materials Science Division, Indira Gandhi Centre for Atomic Research,
Kalpakkam- 603 102. India .

Abstract- We show sensitive dependence on initial conditions and dense set of periodic points imply asymptotic sensitivity, a stronger form of sensitivity, where the deviation happens not just once but infinitely many times. As a consequence it follows that all Devaney chaotic systems (e.g. logistic map) have this asymptotic sensitivity.

Keywords- sensitivity, Devaney's chaos, dense periodic points, asymptotic sensitivity

Sensitive dependence on initial conditions (shortly, sensitivity) is a central concept in the theory chaos and discrete dynamical systems. Roughly speaking, a sensitive system with a sensitivity constant $\delta > 0$ has the following property. Arbitrarily close to the initial point of any specified trajectory, there is a point whose trajectory deviates from that specified one by a distance more than δ , atleast at one instant of time later. The words arbitrarily close emphasizes the fact that even by choosing the initial point closer and closer to that specified initial point one cannot avoid a deviation in distance more than δ . Thus, the arbitrarily small initial separation grows up to more than $\delta > 0$, within a finite time. Hence, in predicting the future an error of magnitude δ is inevitable, however small be the error in the initial condition. This is often referred to, rather crudely, as the divergence of nearby trajectories. The following definition of sensitivity makes this precise.

Discrete dynamical system is a pair (X, T) where X is a metric space and T is a continuous self-map on X . $B_r(x)$ denotes the open ball centered at x and of radius r . The distance between the points x and y of the space X is denoted by $d(x, y)$. The space X is supposed to represent all possible states of a physical system and T models the evolution of the system from the present state x to the next state $T(x)$. Thus the sequence $\{x, T(x), T^2(x), T^3(x), \dots\}$ represents the evolution of the state x and is called the trajectory of x . If $T^k(x) = x$ for some natural number k then x is periodic point. The smallest k for which $T^k(x) = x$ is the order of the periodic point x .

Definition: (sensitivity) The system (X, T) is said to be sensitive if there exists a positive constant δ , called sensitivity constant, such that for all $x \in X$ for all $\epsilon > 0$, there is a natural number $n > 0$ and a pair of points y and z in $B_\epsilon(x)$ such that $d(T^n(z), T^n(y)) > \delta$.

Remark: Note that if $d(T^n(z), T^n(y)) > \delta$ then either $d(T^n(x), T^n(z)) > \delta/2$ or $d(T^n(x), T^n(y)) > \delta/2$. So equivalently with respect to the center of ball $B_\epsilon(x)$, there is a point (y or z) which deviates by $\delta/2$. Some prefer this $\delta/2$ version of the definition.

The question that we ask arises in the following way. The above definition does not preclude the following possibility. All those trajectories, that deviate to a distance more than δ say at some k -th instant, could get closer and closer to that specified trajectory ever after. In other words, all these temporarily deviant trajectories, could asymptotically approach the specified trajectory. In such a system, eventually you will be making lesser and lesser error, in predicting the state of the specified trajectory. Does this happen in the standard sensitive systems like the logistic or a tent map? Are there sensitive maps in which the above scenario of asymptotic convergence of nearby trajectories is impossible? In this note we answer these questions. Any system which has the following property cannot have such asymptotic convergence of nearby trajectories.

Definition: (Asymptotic sensitivity) The dynamical system (X, T) is said to be asymptotically sensitive if there is a $\delta > 0$ such that any ball $B_\epsilon(x)$ (arbitrary $x \in X$ and arbitrary $\epsilon > 0$) contains points y and z such that $d(T^{n_i}(y), T^{n_i}(z)) > \delta$ for infinitely many positive numbers n_i .

Note: Consider the space X of one sided infinite strings from the alphabet $\{0, 1\}$. The shift dynamical system S , acting on this space maps the string $a = a_0a_1a_2\dots$ to $S(a) = a_1a_2a_3\dots$ where a_i is either 1 or 0. The distance between two strings a and b is $d(a, b) = \sum_{i=0}^{\infty} \frac{D(a_i, b_i)}{2^i}$ where D is the discrete metric in the space of two symbols $\{0, 1\}$. Let A be the subspace of X consisting of strings whose terms are eventually all zero i.e. beyond some n th term all the terms are zero. The restriction of T to A is an example of a dynamical system which is sensitive but not asymptotically sensitive [1].

Theorem: If a dynamical system (X, T) is sensitive and has dense set of

periodic points then it is asymptotically sensitive.

Proof: Let the sensitivity constant of the system be δ . The main idea is to choose the diverging pair of points to be periodic points. So one has to show that in any open ball, there exists a pair of periodic point themselves, which separate to a distance more than δ . Since these points are periodic, this separation would occur infinite number of times leading to asymptotic sensitivity. By sensitivity, in any open ball $B_r(x)$ there are points z, y such that

$$d(T^k(z), T^k(y)) > \delta$$

for some integer $k > 1$. Since T^k is continuous, for any open ball $B_\epsilon(T^k(z))$ there is a $\rho > 0$ such that

$$T^k(B_\rho(z)) \subset B_\epsilon(T^k(z))$$

since $U = B_r(x) \cap B_\rho(z)$ is a non-empty open set (contains z), it has a periodic point p (set of periodic points being dense) such that $T^k(p) \in B_\epsilon(T^k(z))$. Similarly there is a periodic point $q \in B_r(x)$ such that $T^k(q) \in B_\epsilon(T^k(y))$. Now we have two periodic points $p, q \in B_r(x)$ such that

$$T^k(p) \in B_\epsilon(T^k(z))$$

and

$$T^k(q) \in B_\epsilon(T^k(y))$$

Since ϵ is arbitrary it is possible to choose the ϵ such that $d(T^k(p), T^k(q)) > \delta$. Let ϵ be chosen such that

$$d(T^k(z), T^k(y)) - 4\epsilon > \delta$$

Then applying triangle inequality to the points $T^k(z), T^k(y)$ and $T^k(p)$ one gets

$$d(T^k(p), T^k(y)) \geq d(T^k(z), T^k(y)) - d(T^k(z), T^k(p))$$

and as $d(T^k(z), T^k(p)) < \epsilon$ it follows

$$d(T^k(p), T^k(y)) \geq d(T^k(z), T^k(y)) - \epsilon$$

Similarly applying triangle inequality to $T^k(p), T^k(q)$ and $T^k(y)$ and using the fact that $d(T^k(q), T^k(y)) < \epsilon$ one gets

$$d(T^k(p), T^k(q)) \geq d(T^k(p), T^k(y)) - \epsilon$$

Combining these one gets

$$d(T^k(p), T^k(q)) \geq d(T^k(z), T^k(y)) - 2\epsilon$$

As $d(T^k(y), T^k(z)) > \delta + 4\epsilon$ by the earlier choice of ϵ it follows

$$d(T^k(p), T^k(q)) > \delta$$

By construction the points p and q are periodic points in the ball $B_r(x)$ and hence $d(T^{k_i}(p), T^{k_i}(q)) > \delta$ for infinitely many positive integers k_i . This proves the proposition.

Du [2], proves amongst others that if the system (X, T) is topologically transitive and has dense set of periodic points then T^k is asymptotically sensitive for every positive k . However, we assume here only sensitivity and dense set of periodic points. Example of a dynamical system which is sensitive and has a dense set periodic points but is not topologically transitive is given in [3].

Since Devaney chaotic systems are sensitive and have dense set of periodic points it follows from the theorem above that they are all asymptotically sensitive as well. For e.g. the well known dynamical systems $([0, 1], 4x(1-x))$ (logistic map) and $([0, 1], 1 - |2x - 1|)$ (tent map) are asymptotic sensitive.

Acknowledgement. The author thanks Prof.V.Kannan and his open neighbourhood for many illuminating discussions and Dr K.P.N. Murthy for his encouragement and support.

References

1. V.Kannan, University of Hyderabad, Personal communication, (2002).
2. Bau sen Du, A dense orbit almost implies sensitivity to initial conditions, Bull. Inst. Math. Acad. Sinica. 26 (1998), 85-94.
3. David Asaaf and Steve Gadbois, Definition of chaos, American mathematical Monthly, (1992) p-865.